

Investigating Redundancy in the Elmo Family and Genetic Pinpointing of Adhesion Defects in *A. thaliana*

Garrison Asper, Class of 2022

For the summer of 2021, I worked to identify plants that contained two mutations within the Elmo family of proteins from a population of *A. thaliana*. The Elmo family consists of five small proteins that are predicted to be localized to the golgi membrane, and serve a function in cellular adhesion. The family was characterized after the discovery of Elmo1 by the Kohorn lab. This protein disrupts cellular adhesion when mutated, and was found using a red dye assay that stains plants with abnormal cellular adhesion. For this project, seed lines containing DNA inserts intended to disrupt each of the remaining four Elmo genes were obtained. The resulting plants were strategically crossed with each other in order to attempt to obtain plants containing two of the possible five mutations in the Elmo family. The double mutant candidates were the progeny of parents that were heterozygous for at least two separate mutations in genes coding for Elmo2 through Elmo5. Candidate plants were screened using a polymerase chain reaction intended to determine whether or not the gene was disrupted. Plants homozygous for the DNA inserts exhibit a similar trend in adhesion defects as the original mutant Elmo1 which was caused by a chemically induced mutation. This gives us confidence that the DNA inserts result in effective disabling of the target Elmo genes and proteins. An Elmo1^{-/-}Elmo2^{-/-} double mutant had already been isolated and studied by the Kohorn lab. This double mutant experienced a disruption in cell adhesion that was much more drastic than either the single Elmo1 mutant or the Elmo2 mutant. This suggests that these two Elmo proteins have redundant functions in the regulation of cellular adhesion.

Only one of the five Elmo genes, Elmo4, has shown to exhibit an extreme adhesion defect such as an Elmo1^{-/-}Elmo2^{-/-} double mutant when disrupted. By obtaining the other combination of double mutants in genes that show weak adhesion defects, the redundancy of the Elmo family of proteins can be studi