

on the market or in

clinical trials contain at least one fluorine atom.<sup>2</sup> Trifluoromethyl groups, three fluorines attached to a singular carbon, are especially common additives, as they can increase the steric bulk of the compound relative to hydrogen groups and provide more conformational control.<sup>3</sup> In order to ensure drug safety and effectiveness, the trifluoromethyl group must be added on enantioselectively

interactions as a remaining capabilities, and the S-1-phenylethylamine side chains help to promote

the cis-conformation of the molecule, which makes the compound form the optimal shape for catalysis.<sup>6</sup>

The three peptoids that were synthesized. The L-alaninamide side chain is in a different position relative to the proposed catalytic site (the 2-picolylamine group on the left of the peptoid) to test if steric or electronic reactions have an impact on the reaction.

All three peptoids were synthesized using solid-phase synthesis. Peptoid 1 was synthesized, analyzed via low resolution liquid chromatography/mass spectrometry (LC-MS) and purified via preparatory high-performance liquid chromatography (HPLC). Purification resulted in a 31% product yield, 50% again in an Purif a pr

1. Chan, P. W. Y.; Yakunin, A. F.; Edwards, E. A.; Pai, E. F., Mapping the Reaction Coordinates of Enzymatic Defluorination. *Journal of the American Chemical Society* , 133 (19), 7461-7468.
2. Böhm, H. J.; Banner, D.; Bendels, S.; Kansy, M.; Kuhn, B.; Müller, K.; Obst-Sander, U.; Stahl, M., Fluorine in Medicinal Chemistry. *ChemBioChem* , 5 (5), 637-643.
3. Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V., Fluorine in Medicinal Chemistry. *Chemical Society Reviews* , 37 (2), 320-330
4. @ji žL "/Li ž7"/K Ub[ žA "/@ji žE " Hf]Zi cfca Yh\mhf]a Yh\mġj`UbY. Bi Wcd\`]W Hf]Zi cfca Yh\mUhf]cb UbX 6YnæbX" ● ž%Ń f&tž\*, ' Ē+' \$"
5. Londoner, R. Investigating the Effect of Side Chains with Hydrogen Bonding Capabilities on Peptoid Catalysts for Enantioselective Trifluoromethylation of 4-Chlorobenzaldehyde. Bowdoin College, Brunswick, Maine, 2020.
6. Wu, C. W.; Sanborn, T. J.; Huang, K.; Zuckermann, R. N.; Barron, A. E., Peptoid Oligomers with -Chiral, Aromatic Side Chains: Sequence Requirements for the Formation of Stable Peptoid Helices. *Journal of the American Chemical Society* , 123 (28), 6778-6784.